


# Pneumococcal Vaccine Timing—For Adults

DO NOT administer PCV13 and PPSV23 at the same visit.

## Age 65 Years or Older

- If PCV13 was given before age 65 years, no additional PCV13 is needed.



## Age 19-64 Years With Underlying Condition(s)

- Prior doses count towards doses recommended below and do not need to be repeated.
- If PPSV23 given previously – wait one year before giving PCV13
  - for group B, wait at least five years before giving a second dose of PPSV23.
- No more than two doses of PPSV23 recommended before 65th birthday and one dose thereafter.

### A. Smoker, Long-term facility resident, or Chronic conditions:

- heart disease (excluding hypertension)
- lung disease (including asthma)
- liver disease (including cirrhosis)
- diabetes
- alcoholism

**PPSV 23**

### B. Immunocompromised (including HIV infection), Chronic renal failure, Nephrotic syndrome, or Asplenia

**PCV 13**

**8 weeks**

**PPSV 23**

**5 years**

**PPSV 23**

### C. CSF leaks or Cochlear implants

**PCV 13**

**8 weeks**

**PPSV 23**



For further details, see: [www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/pneumo.html](http://www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/pneumo.html)

California Department of Public Health, Immunization Branch [www.EZIZ.org](http://www.EZIZ.org)

This publication was supported by Grant Number H23/CCH922507 from the Centers for Disease Control and Prevention (CDC)

IMM-1152 (9/15)