

RUG-IV Systems (Resource Utilization Groups)

		RUG	ADL INDEX
REHABILITATION & EXTENSIVE SERVICES	Ultra High	RUX RUL	11-16 2-10
	Very High	RVX RVL	11-16 2-10
	High	RHX RHL	11-16 2-10
	Medium	RMX RML	11-16 2-10
	Low	RLX	2-16

REHABILITATION ONLY			
720 mins/wk; 2 discipline minimum: one 5+days/wk, one 3+ days/wk; OR Medicare Short Stay = yes with average minutes of 144 or more	Ultra High	RUC RUB RUA	11-16 6-10 0-5
500 mins/wk; 1 discipline minimum: 5+ days/wk; OR Medicare Short Stay = yes with average mins of 100-143	Very High	RVC RVB RVA	11-16 6-10 0-5
325 mins/wk; 1 discipline minimum: 5+ days/wk; OR Medicare Short Stay = yes with average mins of 65-99	High	RHC RHB RHA	11-16 6-10 0-5
150 mins/wk; 5 days across up to 3 disciplines; OR Medicare Short Stay = yes with average minutes of 65-99	Medium	RMC RMB RMA	11-16 6-10 0-5
45 mins/wk; 3 days/wk any combination of 3 disciplines and 6 days of restorative care for 2 services/days; OR Medicare Short Stay = yes with average minutes of 15-29	Low	RLB RLA	11-16 0-10

EXTENSIVE SERVICES			
Residents receiving the following complete clinical care: Tracheostomy care; ventilator/ respirator; isolation for active infectious disease while a resident AND ADL score >=2	Trach care AND ventilator/ respirator	ES3	2-16
	Trach care OR ventilator/ respirator	ES2	2-16
	Isolation for active infectious disease	ES1	2-16

SPECIAL CARE HIGH				
Residents receiving the following complex clinical care or with a following medical condition: Comatose and completely ADL dependent; septicemia; diabetes with daily injections requiring physician order changes on 2 or more days; quadriplegia and ADL score >=5; chronic obstruction pulmonary discarse and shortness of breath when lying flat; fever with pneumonia, or vomiting, or tube feeding, or weight loss; parenteral/IV feedings; respiratory therapy for 7 days; AND ADL>=2 Signs of depression PHQ score >=9.5	Depression Indicators	Yes	HE2	15-16D
		No	HE1	15-16
		Yes	HD2	11-14D
		No	HD1	11-14
		Yes	HC2	6-10D
		No	HC1	6-10
		Yes	HB2	2-5D
		No	HB1	2-5

SPECIAL CARE LOW				
Residents receiving the following complex clinical care or with a following medical condition: Cerebral palsy and ADL score >=5; multiple sclerosis and ADL score >=5; Parkinson's disease and ADL score >=5; feeding tube (calories >=51% or calories = 26-50% and fluid >= 501 cc); ulcers (2 or more stage II or 1 or more stage III or IV pressure ulcers; or 2 or more venous/arterial ulcers; or 1 stage II pressure ulcer and 1 venous/arterial ulcer) with 2 or more skin care treatments; foot infection, respiratory failure and oxygen therapy while a resident; dialysis while a resident; AND ADL score >=2 Signs of depression PHQ score >=9.5	Depression Indicators	Yes	HE2	15-16D
		No	HE1	15-16
		Yes	HD2	11-14D
		No	HD1	11-14
		Yes	HC2	6-10D
		No	HC1	6-10
		Yes	HB2	2-5D
		No	HB1	2-5

CLINICALLY COMPLEX				
Residents with Extensive Services, Special Care High or Special Care low qualifier AND ADL score = 0-1 OR Residents with any of the following clinically complex qualifies: Pneumonia; hemiplegia and ADL score >=5; surgical wounds or open lesions with treatment; burns; chemotherapy while a resident; oxygen while a resident; IV medications while a resident; transfusions while a resident. Signs of depression PHQ score >=9.5	Depression Indicators	Yes	CE2	15-16D
		No	CE1	15-16
		Yes	CD2	11-14D
		No	CD1	11-14
		Yes	CC2	6-10D
		No	CC1	6-10
		Yes	CB2	2-5D
		No	CB1	2-5
		Yes	CA2	0-1 D
		No	CA1	0-1

BEHAVIOR SYMPTOMS & COGNITIVE PERFORMANCE				
Residents having cognitive impairment BIMS score <=9 or CPS >=3 OR Hallucinations of delusions OR Physical or verbal behavioral symptoms toward others; other behavioral symptoms; rejection of care; or wandering exhibited 4+ days over the past 7 days AND ADL score <=5	Nursing Rehab	NR 2+	BB2	2-5
		NR 0-1	BB1	2-5
		NR 2+	BA2	0-1
		NR 0-1	BA1	0-1

REDUCED PHYSICAL FUNCTIONING				
Residents whose needs are primarily for activities of daily living and general supervision. Residents not qualifying for other categories. <u>Nursing Rehab Services:</u> Urinary and/or bowel training program; passive and/or active ROM; amputation prosthesis care training; splint or brace assistance; dressing or grooming training; eating or swallowing training; transfer training; bed mobility and/or walking training; communication training.	Nursing Rehab	NR 2+	PE2	15-16
		NR 0-1	PE1	15-16
		NR 2+	PD2	11-14
		NR 0-1	PD1	11-14
		NR 2+	PC2	6-10
		NR 0-1	PC1	6-10
		NR 2+	PB2	2-5
		NR 0-1	PB1	2-5
NR 2+	PA2	0-1		
NR 0-1	PA1	0-1		