

| Substances: Category and Name | Examples of <i>Commercial</i> and Street Names | DEA Schedule*/ How Administered** | <i>Acute Effects/Health Risks</i> |
|----------------------------------|---|---|--|
| Tobacco | | | |
| Nicotine | Found in cigarettes, cigars, bidis, and smokeless tobacco (snuff, spit tobacco, chew) | Not scheduled/smoked, snorted, chewed | <i>Increased blood pressure and heart rate</i> /chronic lung disease; cardiovascular disease; stroke; cancers of the mouth, pharynx, larynx, esophagus, stomach, pancreas, cervix, kidney, bladder, and acute myeloid leukemia; adverse pregnancy outcomes; addiction |
| Alcohol | | | |
| Alcohol (ethyl alcohol) | Found in liquor, beer, and wine | Not scheduled/swallowed | <i>In low doses, euphoria, mild stimulation, relaxation, lowered inhibitions; in higher doses, drowsiness, slurred speech, nausea, emotional volatility, loss of coordination, visual distortions, impaired memory, sexual dysfunction, loss of consciousness</i> /increased risk of injuries, violence, fetal damage (in pregnant women); depression; neurologic deficits; hypertension; liver and heart disease; addiction; fatal overdose |
| Cannabinoids | | | |
| Marijuana | Blunt, dope, ganja, grass, herb, joint, bud, Mary Jane, pot, reefer, green, trees, smoke, sinsemilla, skunk, weed | I/smoked, swallowed | <i>Euphoria; relaxation; slowed reaction time; distorted sensory perception; impaired balance and coordination; increased heart rate and appetite; impaired learning, memory; anxiety; panic attacks; psychosis</i> /cough; frequent respiratory infections; possible mental health decline; addiction |
| Hashish | Boom, gangster, hash, hash oil, hemp | I/smoked, swallowed | |
| Opioids | | | |
| Heroin | <i>Diacetylmorphine</i> : smack, horse, brown sugar, dope, H, junk, skag, skunk, white horse, China white; cheese (with OTC cold medicine and antihistamine) | I/injected, smoked, snorted | <i>Euphoria; drowsiness; impaired coordination; dizziness; confusion; nausea; sedation; feeling of heaviness in the body; slowed or arrested breathing</i> /constipation; endocarditis; hepatitis; HIV; addiction; fatal overdose |
| Opium | <i>Laudanum, paregoric</i> : big O, black stuff, block, gum, hop | II, III, V/swallowed, smoked | |
| Stimulants | | | |
| Cocaine | <i>Cocaine hydrochloride</i> : blow, bump, C, candy, Charlie, coke, crack, flake, rock, snow, toot | II/snorted, smoked, injected | <i>Increased heart rate, blood pressure, body temperature, metabolism; feelings of exhilaration; increased energy, mental alertness; tremors; reduced appetite; irritability; anxiety; panic; paranoia; violent behavior; psychosis</i> /weight loss; insomnia; cardiac or cardiovascular complications; stroke; seizures; addiction Also, for cocaine —nasal damage from snorting Also, for methamphetamine —severe dental problems |
| Amphetamine | <i>Biphetamine, Dexedrine</i> : bennies, black beauties, crosses, hearts, LA turnaround, speed, truck drivers, uppers | II/swallowed, snorted, smoked, injected | |
| Methamphetamine | <i>Desoxyn</i> : meth, ice, crank, chalk, crystal, fire, glass, go fast, speed | II/swallowed, snorted, smoked, injected | |
| Club Drugs | | | |
| MDMA (methylenedioxyamphetamine) | Ecstasy, Adam, clarity, Eve, lover's speed, peace, uppers | I/swallowed, snorted, injected | MDMA — <i>mild hallucinogenic effects; increased tactile sensitivity, empathic feelings; lowered inhibition; anxiety; chills; sweating; teeth clenching; muscle cramping</i> /sleep disturbances; depression; impaired memory; hyperthermia; addiction Flunitrazepam — <i>sedation; muscle relaxation; confusion; memory loss; dizziness; impaired coordination</i> /addiction GHB — <i>drowsiness; nausea; headache; disorientation; loss of coordination; memory loss</i> /unconsciousness; seizures; coma |
| Flunitrazepam*** | <i>Rohypnol</i> : forget-me pill, Mexican Valium, R2, roach, Roche, roofies, roofinol, rope, rophies | IV/swallowed, snorted | |
| GHB*** | <i>Gamma-hydroxybutyrate</i> : G, Georgia home boy, grievous bodily harm, liquid ecstasy, soap, scoop, goop, liquid X | I/swallowed | |
| Dissociative Drugs | | | |
| Ketamine | <i>Ketalar SV</i> : cat Valium, K, Special K, vitamin K | III/injected, snorted, smoked | <i>Feelings of being separate from one's body and environment; impaired motor function</i> /anxiety; tremors; numbness; memory loss; nausea Also, for ketamine — <i>analgesia; impaired memory; delirium; respiratory depression and arrest; death</i> |
| PCP and analogs | <i>Phencyclidine</i> : angel dust, boat, hog, love boat, peace pill | I, II/swallowed, smoked, injected | Also, for PCP and analogs — <i>analgesia; psychosis; aggression; violence; slurred speech; loss of coordination; hallucinations</i> |
| Salvia divinorum | Salvia, Shepherdess's Herb, Maria Pastora, magic mint, Sally-D | Not scheduled/chewed, swallowed, smoked | Also, for DXM — <i>euphoria; slurred speech; confusion; dizziness; distorted visual perceptions</i> |
| Dextromethorphan (DXM) | Found in some cough and cold medications: Robotripping, Robo, Triple C | Not scheduled/swallowed | |
| Hallucinogens | | | |
| LSD | <i>Lysergic acid diethylamide</i> : acid, blotter, cubes, microdot, yellow sunshine, blue heaven | I/swallowed, absorbed through mouth tissues | <i>Altered states of perception and feeling; hallucinations; nausea</i> Also, for LSD and mescaline — <i>increased body temperature, heart rate, blood pressure; loss of appetite; sweating; sleeplessness; numbness; dizziness; weakness; tremors; impulsive behavior; rapid shifts in emotion</i> |
| Mescaline | Buttons, cactus, mesc, peyote | I/swallowed, smoked | |
| Psilocybin | Magic mushrooms, purple passion, shrooms, little smoke | I/swallowed | Also, for LSD —Flashbacks, Hallucinogen Persisting Perception Disorder Also, for psilocybin — <i>nervousness; paranoia; panic</i> |
| Other Compounds | | | |
| Anabolic steroids | <i>Anadrol, Oxandrin, Durabolin, Depo-Testosterone, Equipoise</i> : roids, juice, gym candy, pumpers | III/injected, swallowed, applied to skin | Steroids — <i>no intoxication effects</i> /hypertension; blood clotting and cholesterol changes; liver cysts; hostility and aggression; acne; in adolescents—premature stoppage of growth; in males—prostate cancer, reduced sperm production, shrunken testicles, breast enlargement; in females—menstrual irregularities, development of beard and other masculine characteristics Inhalants (<i>varies by chemical</i>)— <i>stimulation; loss of inhibition; headache; nausea or vomiting; slurred speech; loss of motor coordination; wheezing</i> /cramps; muscle weakness; depression; memory impairment; damage to cardiovascular and nervous systems; unconsciousness; sudden death |
| Inhalants | <i>Solvents</i> (paint thinners, gasoline, glues); <i>gases</i> (butane, propane, aerosol propellants, nitrous oxide); <i>nitrites</i> (isoamyl, isobutyl, cyclohexyl): laughing gas, poppers, snappers, whippets | Not scheduled/inhaled through nose or mouth | |

| Substances: Category and Name | Examples of <i>Commercial</i> and Street Names | DEA Schedule*/ How Administered** | Acute Effects/Health Risks |
|---------------------------------|--|-----------------------------------|----------------------------|
| Prescription Medications | | | |
| CNS Depressants | For more information on prescription medications, please visit http://www.nida.nih.gov/DrugPages/PrescripDrugsChart.html . | | |
| Stimulants | | | |
| Opioid Pain Relievers | | | |

* Schedule I and II drugs have a high potential for abuse. They require greater storage security and have a quota on manufacturing, among other restrictions. Schedule I drugs are available for research only and have no approved medical use; Schedule II drugs are available only by prescription (unrefillable) and require a form for ordering. Schedule III and IV drugs are available by prescription, may have five refills in 6 months, and may be ordered orally. Some Schedule V drugs are available over the counter.

** Some of the health risks are directly related to the route of drug administration. For example, injection drug use can increase the risk of infection through needle contamination with staphylococci, HIV, hepatitis, and other organisms.

*** Associated with sexual assaults.



Principles of Drug Addiction Treatment

More than three decades of scientific research show that treatment can help drug-addicted individuals stop drug use, avoid relapse and successfully recover their lives. Based on this research, 13 fundamental principles that characterize effective drug abuse treatment have been developed. These principles are detailed in *NIDA's Principles of Drug Addiction Treatment: A Research-Based Guide*. The guide also describes different types of science-based treatments and provides answers to commonly asked questions.

- Addiction is a complex but treatable disease that affects brain function and behavior.** Drugs alter the brain's structure and how it functions, resulting in changes that persist long after drug use has ceased. This may help explain why abusers are at risk for relapse even after long periods of abstinence.
- No single treatment is appropriate for everyone.** Matching treatment settings, interventions, and services to an individual's particular problems and needs is critical to his or her ultimate success.
- Treatment needs to be readily available.** Because drug-addicted individuals may be uncertain about entering treatment, taking advantage of available services the moment people are ready for treatment is critical. Potential patients can be lost if treatment is not immediately available or readily accessible.
- Effective treatment attends to multiple needs of the individual, not just his or her drug abuse.** To be effective, treatment must address the individual's drug abuse and any associated medical, psychological, social, vocational, and legal problems.
- Remaining in treatment for an adequate period of time is critical.** The appropriate duration for an individual depends on the type and degree of his or her problems and needs. Research indicates that most addicted individuals need at least 3 months in treatment to significantly reduce or stop their drug use and that the best outcomes occur with longer durations of treatment.
- Counseling—individual and/or group—and other behavioral therapies are the most commonly used forms of drug abuse treatment.** Behavioral therapies vary in their focus and may involve addressing a patient's motivations to change, building skills to resist drug use, replacing drug-using activities with constructive and rewarding activities, improving problem-solving skills, and facilitating better interpersonal relationships.
- Medications are an important element of treatment for many patients, especially when combined with counseling and other behavioral therapies.** For example, methadone and buprenorphine are effective in helping individuals addicted to heroin or other opioids stabilize their lives and reduce their illicit drug use. Also, for persons addicted to nicotine, a nicotine replacement product (nicotine patches or gum) or an oral medication (bupropion or varenicline), can be an effective component of treatment when part of a comprehensive behavioral treatment program.
- An individual's treatment and services plan must be assessed continually and modified as necessary to ensure it meets his or her changing needs.** A patient may require varying combinations of services and treatment components during the course of treatment and recovery. In addition to counseling or psychotherapy, a patient may

require medication, medical services, family therapy, parenting instruction, vocational rehabilitation and/or social and legal services. For many patients, a continuing care approach provides the best results, with treatment intensity varying according to a person's changing needs.

- Many drug-addicted individuals also have other mental disorders.** Because drug abuse and addiction—both of which are mental disorders—often co-occur with other mental illnesses, patients presenting with one condition should be assessed for the other(s). And when these problems co-occur, treatment should address both (or all), including the use of medications as appropriate.
- Medically assisted detoxification is only the first stage of addiction treatment and by itself does little to change long-term drug abuse.** Although medically assisted detoxification can safely manage the acute physical symptoms of withdrawal, detoxification alone is rarely sufficient to help addicted individuals achieve long-term abstinence. Thus, patients should be encouraged to continue drug treatment following detoxification.
- Treatment does not need to be voluntary to be effective.** Sanctions or enticements from family, employment settings, and/or the criminal justice system can significantly increase treatment entry, retention rates, and the ultimate success of drug treatment interventions.
- Drug use during treatment must be monitored continuously, as lapses during treatment do occur.** Knowing their drug use is being monitored can be a powerful incentive for patients and can help them withstand urges to use drugs. Monitoring also provides an early indication of a return to drug use, signaling a possible need to adjust an individual's treatment plan to better meet his or her needs.
- Treatment programs should assess patients for the presence of HIV/AIDS, hepatitis B and C, tuberculosis, and other infectious diseases, as well as provide targeted risk-reduction counseling to help patients modify or change behaviors that place them at risk of contracting or spreading infectious diseases.** Targeted counseling specifically focused on reducing infectious disease risk can help patients further reduce or avoid substance-related and other high-risk behaviors. Treatment providers should encourage and support HIV screening and inform patients that highly active antiretroviral therapy (HAART) has proven effective in combating HIV, including among drug-abusing populations.



NATIONAL INSTITUTE ON DRUG ABUSE

 RESEARCH DISSEMINATION CENTER

**Order NIDA publications from DrugPubs:
 1-877-643-2644 or 1-240-645-0228 (TTY/TDD)**

This chart may be reprinted. Citation of the source is appreciated.